

PIERI'S FORMULA VIA EXPLICIT RATIONAL EQUIVALENCE

FRANK SOTTILE

ABSTRACT. Pieri's formula describes the intersection product of a Schubert cycle by a special Schubert cycle on a Grassmannian. Classically, it is proven by calculating an appropriate triple intersection of Schubert varieties. We present a new geometric proof, exhibiting an explicit chain of rational equivalences from a suitable sum of distinct Schubert varieties to the intersection of a Schubert variety with a special Schubert variety. The geometry of these rational equivalences indicate a link to a combinatorial proof of Pieri's formula using Schensted insertion.

1. INTRODUCTION

Pieri's formula asserts that the product of a Schubert class and a special Schubert class is a sum of certain other Schubert classes, each with coefficient 1. This determines the multiplicative structure of the Chow ring of a Grassmann variety. Pieri's formula also arises in algebraic combinatorics and representation theory [F], and has independent proofs in each context. One such proof [F] in combinatorics involves Schensted insertion [Sc]. Its geometric proof (for example, in Hodge and Pedoe [HP]) involves studying an intersection of three Schubert varieties and invoking Poincaré duality to obtain the desired sum. We present a new geometric proof of this formula, explicitly describing a sequence of deformations (inducing rational equivalence) that transform a general intersection of a Schubert variety with a special Schubert variety into a sum of distinct Schubert varieties. The geometry of these deformations is quite interesting and their form parallels the combinatorial proof of Pieri's formula using Schensted insertion.

Let $\mathbf{G}_m V$ be the Grassmannian of m -dimensional subspaces of an n -dimensional vector space V . A decreasing sequence α of length m , ($n \geq \alpha_1 > \cdots > \alpha_m \geq 1$), and a complete flag $F_\bullet$ in V together determine a Schubert subvariety $\Omega_\alpha F_\bullet$ of $\mathbf{G}_m V$. Special Schubert varieties Ω_L are those Schubert varieties given by the single condition that an m -plane intersect a given linear subspace L non-trivially. For any subscheme X of $\mathbf{G}_m V$, let $[X]$ be the cycle class of X in the Chow ring of $\mathbf{G}_m V$. Pieri's formula asserts

$$[\Omega_\alpha F_\bullet] \cdot [\Omega_L] = \sum [\Omega_\gamma F_\bullet], \quad (1)$$

the sum over all sequences γ with $\gamma_1 \geq \alpha_1 > \gamma_2 \geq \cdots > \gamma_m \geq \alpha_m$ where $\sum \gamma_i - \alpha_i$ is equal to the codimension b of Ω_L . Let $\alpha * b$ denote this set of sequences and let $Y_{\alpha,b}$ be the

Date: 8 May 1996.

1991 Mathematics Subject Classification. 14M15.

Key words and phrases. Pieri's formula, rational equivalence, Grassmannian, Schensted insertion.

Research supported in part by NSERC grant # OGP0170279.

revised version of alg-geom preprint # 9601006.

cycle $\sum_{\gamma \in \alpha * b} \Omega_\gamma F_\bullet$. Let $\text{Chow } \mathbf{G}_m V$ be the Chow variety of $\mathbf{G}_m V$ and let $\mathcal{G} \subset \text{Chow } \mathbf{G}_m V$ be the set of cycles $\Omega_\alpha F_\bullet \cap \Omega_L$ for all L of a fixed dimension such that the intersection is generically transverse.

Our proof involves a partial compactification of $\mathcal{G}$ in $\text{Chow } \mathbf{G}_m V$ with $b+1$ rational strata, each an orbit of the Borel subgroup of $GL(V)$ stabilizing $F_\bullet$, hence consisting of isomorphic cycles. The 0th stratum is dense in $\mathcal{G}$ and cycles in the i th stratum have components X_β indexed by $\beta \in \alpha * i$, where the component X_β is a subvariety of $\Omega_\beta F_\bullet$. Passing from one stratum to the next, each component X_β deforms into some components of cycles in the next stratum. The ‘history’ of each component $\Omega_\gamma F_\bullet$ of $Y_{\alpha,b}$ through this process gives a chain in the Bruhat order of Schubert varieties, recording which component at each stage gave rise to $\Omega_\gamma F_\bullet$. This leads to the following interpretation of Pieri’s formula: The sum in (1) is over a certain set of chains in the Bruhat order which begin at α , the chain with endpoint γ recording the history of the cycle $\Omega_\gamma F_\bullet$ in the sequence of deformations. In §4, we show how this is similar to a combinatorial proof of Pieri’s formula based on Schensted insertion.

In [Sol], these deformations were constructed in the special case of $\mathbf{G}_2 V$ and applied to obtain a completely geometric understanding of intersections of Schubert subvarieties of $\mathbf{G}_2 V$ in terms of explicit, multiplicity-free deformations. This paper began as an effort to find similar constructions for other Grassmannians, whose geometry is considerably more complicated than that of $\mathbf{G}_2 V$.

This proof of Pieri’s formula is an initial step towards understanding the structure of rational equivalence on these Grassmann varieties in terms of the combinatorics of the Bruhat order of the Schubert cellular decomposition. A chain in the Bruhat order is a standard skew tableau [F]. Thus the Littlewood-Richardson rule for multiplying two Schubert classes has an interpretation as a sum over certain chains in the Bruhat order. A (as yet unknown) geometric proof of the Littlewood-Richardson rule for Grassmannians should provide an explanation for this, similar to what we give for Pieri’s formula.

In fact, we believe that all Schubert-type product formulas for any Grassmannian or flag variety X of any reductive group will eventually be understood in terms of related combinatorics on the Bruhat order on Schubert subvarieties on X , perhaps with additional data giving rise to multiplicities. Some of this picture is already known: Both Chevalley’s [C] formula for multiplication by hypersurface Schubert classes and the Pieri-type formulas of Boe-Hiller [HB] and Pragacz-Ratajski [PR] for Lagrangian and orthogonal Grassmannians are similar to the form of Pieri’s formula (1), but each has multiplicities depending upon root system data. Formulas for multiplying arbitrary Schubert classes in maximal Lagrangian or orthogonal Grassmannians ([P],[St]) are similar to the Littlewood-Richardson formula, using combinatorics of the lattice of shifted Young diagrams, the Bruhat order of these varieties. Known formulas for products in the ordinary flag manifold may also be interpreted in terms of chains in the Bruhat order ([So2],[BS]).

It is only in characteristic zero that general subvarieties of a Grassmannian intersect generically transversally. Kleiman [K] proves this in characteristic zero and gives a counterexample in positive characteristic. In §2, we work over an arbitrary field and give a precise determination (Theorem 2.3) of when a special Schubert variety meets a fixed

Schubert variety generically transversally, and describe the components of such an intersection. The geometry of these components is interesting: while not an intersection of Schubert varieties, each component is ‘birationally fibred’ over such an intersection, with Schubert variety fibres. Such cycles are the key to our proof of Pieri’s formula in §3; they are the components of the intermediate cycles in the deformations used to establish Pieri’s formula.

2. GEOMETRY OF PIERI-TYPE INTERSECTIONS

2.1. Preliminaries. Let k be a fixed, but arbitrary, field and $m \leq n$ positive integers. Let $V \simeq k^n$ be an n -dimensional vector space over k and $\mathbf{G}_m V$ be the Grassmannian of m -planes in V . A *complete flag* $F_\bullet$ in V is a sequence of subspaces

$$0 = F_{n+1} \subset F_n \subset \cdots \subset F_2 \subset F_1 = V$$

of V where $\dim F_j = n + 1 - j$. Let $\langle S \rangle$ denote the linear span of a subset S of V . We let $\binom{[n]}{m}$ be the set of all m -element subsets of $[n] := \{1, 2, \dots, n\}$, considered as decreasing sequences α of length m : $n \geq \alpha_1 > \alpha_2 > \cdots > \alpha_m \geq 1$. A complete flag $F_\bullet$ and a sequence $\alpha \in \binom{[n]}{m}$ together determine a Schubert (sub)variety of $\mathbf{G}_m V$,

$$\Omega_\alpha F_\bullet := \{H \in \mathbf{G}_m V \mid \dim H \cap F_{\alpha_j} \geq j, 1 \leq j \leq m\}.$$

This variety has codimension $|\alpha| := \sum \alpha_i - i$. A *special Schubert variety* is the subvariety of all m -planes H which have a nontrivial intersection with a single subspace F_{m+s} in the flag, $\Omega_{m+s, m-1, \dots, 2, 1} F_\bullet$. We use a compact notation for special Schubert varieties. Let $L := F_{m+s}$, a subspace of dimension $n + 1 - m - s$, and define

$$\Omega_L := \Omega_{m+s, m-1, \dots, 2, 1} F_\bullet.$$

Two subvarieties meet *generically transversally* if they intersect transversally along a dense subset of every component of their intersection. They meet *improperly* if the codimension of their (non-empty) intersection is less than the sum of their codimensions. A subspace L meets a flag $F_\bullet$ *properly* if it meets each subspace F_i properly.

To simplify some assertions and formulae, we adopt the convention that if γ is a decreasing sequence of length m with $\gamma_1 > n$, then $\Omega_\gamma F_\bullet = \emptyset$. Similarly, if the dimension of a subspace is asserted to be negative, we intend that subspace to be $\{0\}$. Also, $\dim\{0\} = -\infty$.

Let $\alpha \in \binom{[n]}{m}$ and r be a positive integer. Define $\alpha * r \subset \binom{[n]}{m}$ to be the set of those $\beta \in \binom{[n]}{m}$ with $\beta_1 \geq \alpha_1 > \beta_2 \geq \cdots > \beta_m \geq \alpha_m$ and $|\beta| = |\alpha| + r$. If $\beta \in \alpha * r$, define $j(\alpha, \beta)$ to be the first index i where β_i differs from α_i , $\min\{i \mid \beta_i > \alpha_i\}$. For $1 \leq j \leq m$, let δ^j be the Kroenecker delta, the sequence with a 1 in the j th position and 0’s elsewhere.

2.2. The cycle $X_\beta(j, F_\bullet, L)$. Central to the geometry of Pieri-type intersections are the components, $X_\beta(j, F_\bullet, L)$, of reducible intersections. These subvarieties are also components of cycles intermediate in deformations establishing Pieri’s formula. Let $\beta \in \binom{[n]}{m}$, $1 \leq j \leq m$ be an integer, $F_\bullet$ a flag, and L a linear subspace in V . Define

$$X_\beta(j, F_\bullet, L) := \{H \in \Omega_\beta F_\bullet \mid \dim H \cap F_{\beta_j} \cap L \geq 1\},$$

a subvariety of $\Omega_\beta F_\bullet \cap \Omega_L$.

The following theorem gives precise conditions on L and $F_\bullet$ which determine whether $\Omega_\alpha F_\bullet \cap \Omega_L$ is improper, generically transverse, or irreducible. Moreover, it computes the components of the intersection in the crucial case of a generically transverse intersection with the maximal number of irreducible components.

2.3. Theorem. *Let $\alpha \in \binom{[n]}{m}$, $s > 0$, $F_\bullet$ be a complete flag in V , and $L \in \mathbf{G}_{n+1-m-s}V$.*

- (1) *Suppose $\dim F_{\alpha_j} \cap L > n + 2 - \alpha_j - j - s$ and $F_{\alpha_j} \cap L \neq \{0\}$, for some $1 \leq j \leq m$. Then $\Omega_\alpha F_\bullet \cap \Omega_L$ is improper. Otherwise, it is generically transverse.*
- (2) *Suppose $\dim F_{\alpha_j} \cap L = n + 2 - \alpha_j - j - s$ for each $1 \leq j \leq m$. Let $M_\bullet$ be any flag satisfying $M_{\alpha_j} = F_{\alpha_j}$ and $M_{\alpha_{j+1}} \supset \langle F_{\alpha_{j-1}}, F_{\alpha_j} \cap L \rangle$, for $1 \leq j \leq m$. Then $\Omega_\alpha F_\bullet$ meets Ω_L generically transversally, and*

$$\Omega_\alpha F_\bullet \cap \Omega_L = \sum_{\beta \in \alpha * 1} X_\beta(j(\alpha, \beta), M_\bullet, L).$$

- (3) *Suppose $\dim F_{\alpha_j} \cap L < n + 2 - \alpha_j - j - s$ for each $1 \leq j < m$ and F_{α_m} meets L properly, so that $\dim F_{\alpha_m} \cap L = n + 2 - \alpha_m - m - s$. Then $\Omega_\alpha F_\bullet \cap \Omega_L$ is irreducible.*

Note that $n + 2 - \alpha_j - j - s$, the critical dimension for $F_{\alpha_j} \cap L$ in this theorem, exceeds the expected dimension of $n + 2 - \alpha_j - m - s$ by $m - j$. Thus, it is not necessary for $F_\bullet$ and L to meet properly for $\Omega_\alpha F_\bullet \cap \Omega_L$ to be generically transverse or even irreducible. However, it is necessary that F_{α_m} and L meet properly. We also see that, as the relative position of $F_\bullet$ and L becomes more degenerate, the intersection $\Omega_\alpha F_\bullet \cap \Omega_L$ ‘branches’ into components, one for each j such that $\dim F_{\alpha_j} \cap L = n + 2 - \alpha_j - j - s$, and it will attain excess intersection if $\dim F_{\alpha_j} \cap L > n + 2 - \alpha_j - j - s$, for even one j .

2.4. Remark. In the situation of Theorem 2.3(2), if $\beta \in \alpha * 1$ and $j(\alpha, \beta) = 1$, then $\beta = \alpha + \delta^1$. Suppose further that $M_{\alpha_1} \cap L = M_{\alpha_1+s}$. Then

$$X_\beta(1, M_\bullet, L) = \Omega_{\alpha+s\delta^1} M_\bullet = \Omega_{\beta+(s-1)\delta^1} M_\bullet,$$

so we have

$$\Omega_\alpha F_\bullet \cap \Omega_L = \sum_{\substack{\beta \in \alpha * 1 \\ j(\alpha, \beta) = 1}} \Omega_{\beta+(s-1)\delta^1} M_\bullet + \sum_{\substack{\beta \in \alpha * 1 \\ j(\alpha, \beta) > 1}} X_\beta(j(\alpha, \beta), M_\bullet, L).$$

We prove Theorem 2.3 in §2.9. First, we study the varieties $X_\beta(j, F_\bullet, L)$. Let $\beta \in \binom{[n]}{m}$, $F_\bullet$ be a complete flag, and $1 \leq j \leq m$ an integer. The map from $\Omega_\beta F_\bullet$ to $\mathbf{G}_j F_{\beta_j}$ given by $H \mapsto H \cap F_{\beta_j}$ is only defined on the dense locus in $\Omega_\beta F_\bullet$ of those H where $\dim H \cap F_{\beta_j} = j$. Resolving the ambiguity of this map gives the variety

$$\widetilde{\Omega}_\beta^j F_\bullet := \{(H, K) \in \Omega_\beta F_\bullet \times \mathbf{G}_j F_{\beta_j} \mid K \subset H \text{ and } \dim K \cap F_{\beta_i} \geq i, 1 \leq i \leq j\}.$$

In Lemma 2.5, we show that the projection to $\mathbf{G}_j F_{\beta_j}$ realizes $\widetilde{\Omega}_\beta^j F_\bullet$ as a fibre bundle with base and fibres themselves Schubert varieties. The following definitions are needed to describe the base and fibres. Let p be the first projection and π the second. For $K \subset V$, let $F_\bullet/K$ be the image of the flag $F_\bullet$ in V/K . Let $F_\bullet|_{\beta_j}$ be the flag

$$F_{\beta_j} \supset F_{\beta_{j+1}} \supset \cdots \supset F_n$$

and $\beta|_j \in \binom{[n+1-\beta_j]}{j}$ the sequence

$$\beta_1 - \beta_j + 1 > \cdots > \beta_{j-1} - \beta_j + 1 > 1 = (\beta|_j)_j.$$

Unraveling this definition shows $(F|_{\beta_j})_{(\beta|_j)_i} = F_{\beta_i}$, for $i \leq j$.

2.5. Lemma. *Let $\beta \in \binom{[n]}{m}$, $F_\bullet$ be a flag, and $1 \leq j \leq m$. Then p is an isomorphism over the dense subset $\{H \in \Omega_\beta F_\bullet \mid \dim H \cap F_{\beta_j} = j\}$. Also, π exhibits $\widetilde{\Omega}_\beta^j F_\bullet$ as a fibre bundle with base $\Omega_{\beta|_j} F_\bullet|_{\beta_j}$ whose fibre over $K \in \Omega_{\beta|_j} F_\bullet|_{\beta_j}$ is the Schubert variety $\Omega_{\beta_{j+1} \dots \beta_m} F_\bullet / K \subset \mathbf{G}_{m-j} V / K$. Moreover, each fibre of π meets the locus where p is an isomorphism.*

Proof: We describe the fibres of π . Note that Schubert varieties have a dual description:

$$H \in \Omega_\beta F_\bullet \iff \dim \frac{H}{H \cap F_{\beta_i}} \leq m - i, \text{ for } 1 \leq i \leq m.$$

If $K \in \Omega_{\beta|_j} F_\bullet|_{\beta_j}$, then $K \subset F_{\beta_j} \subset F_{\beta_i}$, for $i > j$. Thus $(F_\bullet / K)_{\beta_i} = F_{\beta_i} / K$, for $i > j$. Hence, if H is in the fibre over K , then $H \in \Omega_\beta F_\bullet$ and $K \subset H$, so

$$\dim \frac{H/K}{H/K \cap (F_\bullet / K)_{\beta_i}} = \dim \frac{H}{H \cap F_{\beta_i}} \leq m - i, \text{ for } j < i \leq m.$$

Thus $H/K \in \Omega_{\beta_{j+1} \dots \beta_m} F_\bullet / K$. The reverse implication is similar and the remaining assertions follow easily from the definitions. $\blacksquare$

Reformulating the definition of $X_\beta(j, F_\bullet, L)$ in these terms gives a useful characterization.

2.6. Corollary. $X_\beta(j, F_\bullet, L) = p(\pi^{-1}(\Omega_{\beta|_j} F_\bullet|_{\beta_j} \cap \Omega_{F_{\beta_j} \cap L}))$.

Since the fibres of π meet the locus where p is an isomorphism, the map

$$p : \pi^{-1}(\Omega_{\beta|_j} F_\bullet|_{\beta_j} \cap \Omega_{F_{\beta_j} \cap L}) \longrightarrow X_\beta(j, F_\bullet, L)$$

is proper and birational. Thus, while $X_\beta(j, F_\bullet, L)$ is neither a Schubert variety nor an intersection of Schubert varieties, it is ‘birationally fibred’ over an intersection of Schubert varieties with Schubert variety fibres, and hence is intermediate between these extremes.

2.7. Tangent spaces to Schubert varieties. Let $H \in \mathbf{G}_m V$ and $K \in \mathbf{G}_{n-m} V$ be complementary subspaces, so $H \cap K = \{0\}$. The open set $U \subset \mathbf{G}_m V$ of those H' with $H' \cap K = \{0\}$ is identified with $\text{Hom}(H, K)$ by $\phi \in \text{Hom}(H, K) \mapsto \Gamma_\phi$, the graph of ϕ in $H \oplus K = V$. This shows the tangent space of $\mathbf{G}_m V$ at H , $T_H \mathbf{G}_m V$, is equal to $\text{Hom}(H, V/H)$, since K is canonically isomorphic to V/H . The intersection of a Schubert variety $\Omega_\alpha F_\bullet$ containing H with this open set U can be used to determine whether $\Omega_\alpha F_\bullet$ is smooth at H and its tangent space at H . This gives the following description: If $H \in \mathbf{G}_m V$ and $\dim H \cap F_{\alpha_j} = j$ for $1 \leq j \leq m$, then $\Omega_\alpha F_\bullet$ is smooth at H and

$$T_H \Omega_\alpha F_\bullet = \{\phi \in \text{Hom}(H, V/H) \mid \phi(H \cap F_{\alpha_j}) \subset (F_{\alpha_j} + H)/H, 1 \leq j \leq m\}.$$

Similarly, if $H \in \mathbf{G}_m V$, $L \in \mathbf{G}_{n+1-m-s} V$, and $\dim H \cap L = 1$, then Ω_L is smooth at H and the tangent space of Ω_L at H is

$$T_H \Omega_L = \{\phi \in \text{Hom}(H, V/H) \mid \phi(H \cap L) \subset (L + H)/H\}.$$

Let P be the subgroup of $GL(V)$ stabilizing the partial flag $F_{\alpha_1} \subset F_{\alpha_2} \subset \cdots \subset F_{\alpha_m}$. The orbit $P \cdot L'$ consists of those L with $\dim F_{\alpha_j} \cap L = \dim F_{\alpha_j} \cap L'$ for $1 \leq j \leq m$. Similarly, $L \in \overline{P \cdot L'}$ if $\dim F_{\alpha_j} \cap L \geq \dim F_{\alpha_j} \cap L'$ for $1 \leq j \leq m$. If $P \cdot L = P \cdot L'$, then $\Omega_\alpha F \cdot \bigcap \Omega_L \simeq \Omega_\alpha F \cdot \bigcap \Omega_{L'}$. Thus P -orbits on $\mathbf{G}_{n+1-m-s}V$ determine the isomorphism type of Pieri-type intersections.

2.8. Lemma. *Suppose that $L, L' \in \mathbf{G}_{n+1-m-s}V$ with $L \in \overline{P \cdot L'}$. Then*

- (1) $\dim \Omega_\alpha F \cdot \bigcap \Omega_L \geq \dim \Omega_\alpha F \cdot \bigcap \Omega_{L'}$.
- (2) *If $\Omega_\alpha F \cdot \bigcap \Omega_L$ is generically transverse, then $\Omega_\alpha F \cdot \bigcap \Omega_{L'}$ is generically transverse.*
- (3) *If $\Omega_\alpha F \cdot \bigcap \Omega_L$ is generically transverse and irreducible, then $\Omega_\alpha F \cdot \bigcap \Omega_{L'}$ is generically transverse and irreducible.*

Proof: Let $\psi : \mathbf{P}^1 \rightarrow \overline{P \cdot L'}$ be a map with $\psi(0) = L$ and $\psi(\mathbf{P}^1) \cap (P \cdot L') \neq \emptyset$. Then $\Omega_\alpha F \cdot \bigcap \Omega_{\psi(t)}$ is isomorphic to $\Omega_\alpha F \cdot \bigcap \Omega_{L'}$, for any $t \in \psi^{-1}(P \cdot L')$. The lemma follows by considering the subvariety of $\mathbf{P}^1 \times \mathbf{G}_m V$ whose fibre over $t \in \mathbf{P}^1$ is $\Omega_\alpha F \cdot \bigcap \Omega_{\psi(t)}$. ■

2.9. Proof of Theorem 2.3: Let $\alpha \in \binom{[n]}{m}$, $s > 0$, $F_\bullet$ be a complete flag, and $L \in \mathbf{G}_{n+1-m-s}V$. The conditions on L in statement (2), that $\dim F_{\alpha_j} \cap L = n + 2 - \alpha_j - j - s$ for each j , determine a P -orbit, which is the closure of any P -orbit $P \cdot L'$, where $\dim F_{\alpha_j} \cap L' \leq n + 2 - \alpha_j - j - s$ for each j . Thus (2) and Lemma 2.8(2) together imply that if $\dim F_{\alpha_j} \cap L \leq n + 2 - \alpha_j - j - s$ for each j , then $\Omega_\alpha F \cdot \bigcap \Omega_L$ is generically transverse, proving the second part of (1).

For the first part of (1), suppose $\dim F_{\alpha_j} \cap L > n + 2 - \alpha_j - j - s$ and let $L' := F_{\alpha_j} \cap L \neq \{0\}$. Then L' has codimension at most $j + s - 1$ in F_{α_j} . Hence $\Omega_{\alpha|_j} F \cdot |_{\alpha_j} \bigcap \Omega_{L'} \neq \emptyset$ and so has codimension in $\Omega_{\alpha|_j} F \cdot |_{\alpha_j}$ at most that of $\Omega_{L'}$ in $\mathbf{G}_j F_{\alpha_j}$, which is at most $s - 1$. Thus

$$X_\alpha(j, F_\bullet, L) = p(\pi^{-1}(\Omega_{\alpha|_j} F \cdot |_{\alpha_j} \bigcap \Omega_{L'}))$$

which has codimension less than s in $\Omega_\alpha F \cdot \bigcap \Omega_L = p(\pi^{-1}(\Omega_{\alpha|_j} F \cdot |_{\alpha_j}))$. Hence $\Omega_\alpha F \cdot \bigcap \Omega_L$ is improper, as $X_\alpha(j, F_\bullet, L) \subset \Omega_\alpha F \cdot \bigcap \Omega_L$, proving (1).

Before proceeding with the rest of the proof, we make a computation. Suppose $\dim F_{\alpha_j} \cap L \leq n + 2 - \alpha_j - j - s$ for $1 \leq j \leq m$ and $F_{\alpha_m} \cap L \not\subset F_{\alpha_{m-1}}$. Then there exists $H \in \Omega_\alpha F \cdot \bigcap \Omega_L$ with $\dim H \cap F_{\alpha_j} = j$ for $1 \leq j \leq m$, $\dim H \cap L = 1$, and $H \cap L \not\subset F_{\alpha_{m-1}}$. Inductively choose linearly independent vectors $f_j \in F_{\alpha_j}$ for $1 \leq j \leq m$ as follows. Let $f_1 \in F_{\alpha_1} - \{0\}$. Then for $1 < j < m$ suppose that $f_1, \dots, f_{j-1}$ have been chosen. Since $\dim F_{\alpha_j}$ exceeds

$$\dim F_{\alpha_j} \cap \langle L, f_1, \dots, f_{j-1} \rangle \leq n + 2 - \alpha_j - j - s + (j - 1) = n + 1 - \alpha_j - s$$

and $F_{\alpha_{j-1}} \not\subset F_{\alpha_j}$, we can select a vector f_j in

$$F_{\alpha_j} - \langle L, f_1, \dots, f_{j-1} \rangle - F_{\alpha_{j-1}}.$$

Let $f_m \in F_m \cap L - F_{\alpha_{m-1}}$, and let $H := \langle f_1, \dots, f_m \rangle$. Then $H \in \Omega_\alpha F \cdot \bigcap \Omega_L$, $\dim H \cap F_{\alpha_j} = j$ for $1 \leq j \leq m$, $\dim H \cap L = 1$, and $H \cap L \not\subset F_{\alpha_{m-1}}$. Let X_m° be the set of all such H . For $H \in X_m^\circ$,

$$T_H \Omega_\alpha F \cdot \bigcap T_H \Omega_L = \{ \phi \in T_H \Omega_\alpha F \cdot | \phi(H \cap L) \subset (F_{\alpha_m} \cap L + H)/H \}.$$

This has codimension in $T_H\Omega_\alpha F_\bullet$ equal to $\dim(F_{\alpha_m} + H) - \dim(F_{\alpha_m} \cap L + H) = s$. Thus $\Omega_\alpha F_\bullet$ and Ω_L meet transversally along X_m° .

We show (2). Suppose $\dim F_{\alpha_j} \cap L = n + 2 - \alpha_j - s$ for each $1 \leq j \leq m$. Let $M_\bullet$ be any flag satisfying $M_{\alpha_j} = F_{\alpha_j}$ and $M_{\alpha_{j+1}} \supset \langle F_{\alpha_{j-1}}, F_{\alpha_j} \cap L \rangle$, for $1 \leq j \leq m$. Let $H \in \Omega_\alpha F_\bullet \cap \Omega_L$. Then there is some $1 \leq j \leq m$ with $H \cap L \cap F_{\alpha_j} \not\subset F_{\alpha_{j-1}}$. Since $\dim H \cap F_{\alpha_{j-1}} \geq j - 1$, we have $\dim H \cap \langle F_{\alpha_{j-1}}, F_{\alpha_j} \cap L \rangle \geq j$ and so $\dim H \cap M_{\alpha_{j+1}} \geq j$. Thus $H \in \Omega_{\alpha+\delta^j} M_\bullet$, if $\alpha + \delta^j \in \binom{[n]}{m}$. But this is the case, as $\alpha_j + 1 < \alpha_{j-1}$, for otherwise dimensional considerations imply that $L \cap F_{\alpha_j} = L \cap F_{\alpha_{j-1}} \subset F_{\alpha_{j-1}}$. Let $\beta := \alpha + \delta^j \in \alpha * 1$. Then $j(\alpha, \beta) = j$ and $H \in X_\beta(j(\alpha, \beta), M_\bullet, L)$, since $H \in \Omega_\beta M_\bullet$ and $\dim H \cap L \cap M_{\beta_j} \geq 1$. Conversely, if $\beta \in \alpha * 1$, then $\Omega_\beta M_\bullet \subset \Omega_\alpha F_\bullet$, so $X_\beta(j(\alpha, \beta), M_\bullet, L) \subset \Omega_\alpha F_\bullet \cap \Omega_L$. This shows

$$\Omega_\alpha F_\bullet \cap \Omega_L = \sum_{\beta \in \alpha * 1} X_\beta(j(\alpha, \beta), M_\bullet, L).$$

We claim this intersection is generically transverse. Let $\beta \in \alpha * 1$ and $j := j(\alpha, \beta)$. Then $X_\beta(j, M_\bullet, L)$ has an open subset X_j° consisting of those H with $\dim H \cap F_{\alpha_i} = i$ for $1 \leq i \leq m$, $\dim H \cap L = 1$, and $H \cap L \subset F_{\alpha_j}$ but $H \cap L \not\subset F_{\alpha_{j-1}}$. As with X_m° above, X_j° is nonempty, so it is a dense open subset of $X_\beta(j, M_\bullet, L)$. For $H \in X_j^\circ$,

$$T_H\Omega_\alpha F_\bullet \cap T_H\Omega_L = \{\phi \in T_H\Omega_\alpha F_\bullet \mid \phi(H \cap L) \subset (L \cap F_{\alpha_j} + H)/H\}.$$

Since $\dim(F_{\alpha_j} + H) - \dim(L \cap F_{\alpha_j} + H) = s$, this has codimension s in $T_H\Omega_\alpha F_\bullet$, showing that $\Omega_\alpha F_\bullet$ and Ω_L meet transversally along X_j° , a dense subset of $X_\beta(j(\alpha, \beta), M_\bullet, L)$.

By Lemma 2.8(3), it suffices to prove a special case of (3):

(3)' *If F_{α_m} meets L properly, and for $1 \leq j < m$, $\dim F_{\alpha_j} \cap L = n + 2 - \alpha_j - j - (s + 1)$, then $\Omega_\alpha F_\bullet \cap \Omega_L$ is irreducible.*

These conditions imply $F_{\alpha_m} \cap L \not\subset F_{\alpha_{m-1}}$. In the notation of §2.4, let $L' := F_{\alpha_{m-1}} \cap L$, $F'_\bullet := F_\bullet|_{\alpha_{m-1}}$, and $\alpha' := \alpha|_{m-1}$. Consider

$$X_\alpha(m-1, F_\bullet, L) = p(\pi^{-1}(\Omega_{\alpha'} F'_\bullet \cap \Omega_{L'})).$$

For $j \leq m-1$,

$$\begin{aligned} \dim F_{\alpha_j} \cap L' &= n + 2 - \alpha_j - j - (s + 1) \\ &= \dim F_{\alpha_{m-1}} + 2 - \alpha'_j - j - (s + 1), \end{aligned}$$

so L' and $F'_\bullet$ satisfy the conditions of (2) for the pair $\alpha', s + 1$. Thus $\Omega_{\alpha'} F'_\bullet \cap \Omega_{L'}$ is generically transverse, which implies that $X_\alpha(m-1, F_\bullet, L)$ has codimension $s + 1$ in $\Omega_\alpha F_\bullet$ and hence is a proper subvariety of $\Omega_\alpha F_\bullet \cap \Omega_L - X_\alpha(m-1, F_\bullet, L)$. Since X_m° is dense in $\Omega_\alpha F_\bullet \cap \Omega_L - X_\alpha(m-1, F_\bullet, L)$, this completes the proof of (3)'. $\blacksquare$

3. CONSTRUCTION OF EXPLICIT RATIONAL EQUIVALENCES

Theorem 2.3 shows that for L in a dense subset of $\mathbf{G}_{n+1-m-b}V$, the intersection $\Omega_\alpha F_\bullet \cap \Omega_L$ is generically transverse and irreducible. We use Theorem 2.3(2) to study such a cycle as L ‘moves out of’ this set, ultimately deforming it into the cycle $\sum_{\gamma \in \alpha * b} \Omega_\gamma F_\bullet$.

3.1. Families and Chow varieties. Suppose $\Sigma \subset (\mathbf{P}^1 - \{0\}) \times \mathbf{G}_m V$ has equidimensional fibres over $\mathbf{P}^1 - \{0\}$. Then its Zariski closure $\overline{\Sigma}$ in $\mathbf{P}^1 \times \mathbf{G}_m V$ has equidimensional fibres over $\mathbf{P}^1$. Denote the fibre of $\overline{\Sigma}$ over 0 by $\lim_{t \rightarrow 0} \Sigma_t$, where Σ_t is the fibre of Σ over $t \in \mathbf{P}^1 - \{0\}$. The association of a point t of $\mathbf{P}^1$ to the fundamental cycle of the fibre $\overline{\Sigma}_t$ determines a morphism $\mathbf{P}^1 \rightarrow \text{Chow } \mathbf{G}_m V$. Moreover, if Σ is defined over k , then so is the map $\mathbf{P}^1 \rightarrow \text{Chow } \mathbf{G}_m V$ ([Sa], §I.9).

3.2. The cycle $Y_{\alpha,r}(F, L)$. In §2.2, we defined the components $X_\beta(\alpha, F, L)$ of the cycles intermediate between $\Omega_\alpha F \cap \Omega_L$ and $\sum_{\gamma \in \alpha * b} \Omega_\gamma F$. Here, we define those intermediate cycles, $Y_{\alpha,r}(F, L)$, which are parameterized by subspaces L in certain Schubert cells $U_{\alpha,s} F$ of $\mathbf{G}_{n+1-m-s} V$. Let $U_{\alpha,s} F$ be the set of those $L \in G_{n+1-m-s} V$ such that

- (1) $F_{\alpha_1} \cap L = F_{\alpha_1+s}$, and
- (2) $F_{\alpha_j} \cap L = F_{\alpha_j+1} \cap L$, and has dimension $n+2-\alpha_j-j-s$, for $1 \leq j \leq m$.

These conditions are consistent and determine $\dim F_i \cap L$ for $1 \leq i \leq n$. For example,

$$(3.2) \quad \alpha_j < i < \alpha_{j-1} \implies \dim F_i \cap L = \dim F_i + 1 - j - (s-1).$$

Thus $U_{\alpha,s} F$ is a single Schubert cell of $G_{n+1-m-s} V$. Specifically, $U_{\alpha,s} F$ is the dense cell of $\Omega_\beta F$, where $\beta \in \binom{[n]}{n+1-m-s}$ is defined as follows: If $\alpha_1 \leq n+1-s$, then $\beta = [n] - \alpha - \{\alpha_1+1, \dots, \alpha_1+s-1\}$. Otherwise, β is the smallest $n+1-m-s$ integers in $[n] - \alpha$.

For $\beta \in \alpha * r$, recall that $j(\alpha, \beta) = \min\{i \mid \alpha_i < \beta_i\}$. If $L \in U_{\alpha,s} F$, define the cycle

$$Y_{\alpha,r}(F, L) := \sum_{\substack{\beta \in \alpha * r \\ j(\alpha, \beta) = 1}} \Omega_{\beta+(s-1)\delta^1} F + \sum_{\substack{\beta \in \alpha * r \\ j(\alpha, \beta) > 1}} X_\beta(j(\alpha, \beta), F, L).$$

Let $\mathcal{G}_{\alpha,s,r} F \subset \text{Chow } \mathbf{G}_m V$ be the set of these cycles $Y_{\alpha,r}(F, L)$ for $L \in U_{\alpha,s} F$. Since $U_{\alpha,s} F$ is a Schubert cell, $\mathcal{G}_{\alpha,s,r} F$ is an orbit of the Borel subgroup stabilizing F and hence is rational.

3.3. Remark. Suppose $L \in U_{\alpha,s} F$, then by Remark 2.4,

$$\begin{aligned} \Omega_\alpha F \cap \Omega_L &= \sum_{\substack{\beta \in \alpha * 1 \\ j(\alpha, \beta) = 1}} \Omega_{\beta+(s-1)\delta^1} F + \sum_{\substack{\beta \in \alpha * 1 \\ j(\alpha, \beta) > 1}} X_\beta(j(\alpha, \beta), F, L) \\ &= Y_{\alpha,1}(F, L). \end{aligned}$$

The following lemma parameterizes our explicit rational equivalences. It is identical to Lemma 6.1 of [Sol].

3.4. Lemma. *Let $l \leq n$ and let M_l be a complete flag in $M \simeq k^n$. Suppose L_∞ is a hyperplane containing M_l but not M_{l-1} . Then there exists a pencil of hyperplanes L_t , for $t \in \mathbf{P}^1$, such that if $t \neq 0$, then L_t contains M_l but not M_{l-1} and, for each $i \leq l-1$, the family of codimension $i+1$ planes induced by $M_i \cap L_t$ for $t \neq 0$ has fibre M_{i+1} over 0.*

Proof: Let $x_1, \dots, x_n$ be a basis of M^* such that $L_\infty = \langle x_{l-1} \rangle^\perp$ and $M_i = \langle x_1, \dots, x_{i-1} \rangle^\perp$. Let $e_1, \dots, e_n$ be a basis for M dual to $x_1, \dots, x_n$ and define

$$L_t := \langle M_l, te_j + e_{j+1} \mid 1 \leq j \leq l-2 \rangle.$$

For $t \neq 0$ and $1 \leq i \leq l-1$, $M_i \cap L_t = \langle M_l, te_j + e_{j+1} \mid i \leq j \leq l-2 \rangle$ and so has dimension $n-i$. The fibre of this family at $t=0$ is $\langle M_l, e_{j+1} \mid i \leq j \leq l-2 \rangle = M_{i+1}$. $\blacksquare$

3.5. Theorem. *Let $\alpha \in \binom{[n]}{m}$, s, r be positive integers and $F_\bullet$ a flag in V . Let $M \in U_{\alpha, s-1} F_\bullet$ and define $M_\bullet$ to be the flag in M consisting of the subspaces in $F_\bullet \cap M$.*

Let $L_\infty \subset M$ be any hyperplane containing F_{α_1+s} but not F_{α_1+s-1} . Suppose L_t is the family of hyperplanes of M given by Lemma 3.4. Then

- (1) *For $t \neq 0$, $L_t \in U_{\alpha, s} F_\bullet$.*
- (2) $\lim_{t \rightarrow 0} Y_{\alpha, r}(F_\bullet, L_t) = Y_{\alpha, r+1}(F_\bullet, M)$.

3.6. Theorem. [Pieri's Formula] *Let $\alpha \in \binom{[n]}{m}$, $F_\bullet$ be a complete flag in V , and $K \in \mathbf{G}_{n+1-m-b} V$ be a subspace which meets $F_\bullet$ properly. Then the cycle $\Omega_\alpha F_\bullet \cap \Omega_K$, a generically transverse intersection, is rationally equivalent to $\sum_{\gamma \in \alpha * b} \Omega_\gamma F_\bullet$. Thus, in the Chow ring $A^* \mathbf{G}_m V$ of $\mathbf{G}_m V$,*

$$[\Omega_\alpha F_\bullet] \cdot [\Omega_K] = \sum_{\gamma \in \alpha * b} [\Omega_\gamma F_\bullet].$$

Moreover, let $\mathcal{G} \subset \text{Chow } \mathbf{G}_m V$ be the set of cycles arising as generically transverse intersections of the form $\Omega_\alpha F_\bullet \cap \Omega_K$ for $K \in \mathbf{G}_{n+1-m-b} V$. Then one may give $b+1$ explicit rational deformations inducing this rational equivalence, where the cycles at the i th stage are of the form $Y_{\alpha, i}(F_\bullet, M)$, with $M \in U_{\alpha, b+1-i} F_\bullet$, and all are within $\mathcal{G}$.

The Borel subgroup B of $GL(V)$ stabilizing $F_\bullet$ also stabilizes $\mathcal{G}$ and the cycle $\sum_{\gamma \in \alpha * b} \Omega_\gamma F_\bullet$ is the only B -stable cycle in this component of the Chow variety. As Hirschowitz [H] observed, this implies $\sum_{\gamma \in \alpha * b} \Omega_\gamma F_\bullet$ is in the Zariski closure of $\mathcal{G}$. Thus Theorem 3.6 is an improvement in that deformations inducing the rational equivalence are given explicitly.

3.7. Proof of Pieri's formula using Theorem 3.5. Let $b > 0$, and $\alpha \in \binom{[n]}{m}$. For $1 \leq i \leq b$, let $U_i := U_{\alpha, b+1-i} F_\bullet$ and $\mathcal{G}_i := \mathcal{G}_{\alpha, b+1-i, i} F_\bullet$. Let $U_0 \subset \mathbf{G}_{n+1-m-b} V$ be the (dense) set of those L which meet F_{α_m} properly and for $1 \leq j < m$, $\dim F_{\alpha_j} \cap L < n+2-\alpha_j-j-b$. By Theorem 2.3, if $L \in \mathbf{G}_{n+1-m-b} V$, then $\Omega_\alpha F_\bullet \cap \Omega_L$ is generically transverse and irreducible if and only if $L \in U_0$. Let $\mathcal{G}_0 \subset \text{Chow } \mathbf{G}_m V$ be the set of cycles $\Omega_\alpha F_\bullet \cap \Omega_L$ for $L \in U_0$.

Let $L \in U_b$ and consider the cycle $Y_{\alpha, b}(F_\bullet, L) \in \mathcal{G}_b$:

$$Y_{\alpha, b}(F_\bullet, L) = \sum_{\substack{\beta \in \alpha * b \\ j(\alpha, \beta)=1}} \Omega_\beta F_\bullet + \sum_{\substack{\beta \in \alpha * b \\ j(\alpha, \beta)>1}} X_\beta(j(\alpha, \beta), F_\bullet, L).$$

We claim $Y_{\alpha, b}(F_\bullet, L) = \sum_{\beta \in \alpha * b} \Omega_\beta F_\bullet$, the cycle $Y_{\alpha, b} F_\bullet$ of the Introduction. It suffices to show $X_\beta(j(\alpha, \beta), F_\bullet, L) = \Omega_\beta F_\bullet$ for $\beta \in \alpha * b$ with $j(\alpha, \beta) > 1$. Suppose $j = j(\alpha, \beta) > 1$, then

$$X_\beta(j, F_\bullet, L) = p(\pi^{-1}(\Omega_{\beta|_j} F_\bullet|_{\beta_j} \cap \Omega_{F_{\beta_j} \cap L})).$$

By formula (3.2), $\dim F_{\beta_j} \cap L = \dim F_{\beta_j} - j + 1$, as $\alpha_j < \beta_j < \alpha_{j-1}$ and $s = 1$. So $\Omega_{F_{\beta_j} \cap L}$ is $\mathbf{G}_j F_{\beta_j}$, since any j -plane in F_{β_j} meets $F_{\beta_j} \cap L$ non-trivially. Thus $X_\beta(j(\alpha, \beta), F_\bullet, L) = \Omega_\beta F_\bullet$, by the definition of p and π in §2.4.

Let $\mathcal{G} \subset \text{Chow } \mathbf{G}_m V$ be the set of all cycles $\Omega_\alpha F \cap \Omega_L$, where $L \in \mathbf{G}_{n+1-m-b} V$ and the intersection is generically transverse. Then by Theorem 2.3 and Remark 3.3, both $\mathcal{G}_0$ and $\mathcal{G}_1$ are subsets of $\mathcal{G}$. Arguing as in the proof of Lemma 2.8 shows $\mathcal{G} \subset \overline{\mathcal{G}_0}$. Theorem 3.5 implies $\mathcal{G}_i \subset \overline{\mathcal{G}_{i-1}}$ for $2 \leq i \leq b$, so in particular, $Y_{\alpha,b} F \in \mathcal{G}_b \subset \overline{\mathcal{G}}$. Since $\mathcal{G}_0$ and hence $\overline{\mathcal{G}}$ is rational, $Y_{\alpha,b} F$ is rationally equivalent to any cycle in $\mathcal{G}$, including $\Omega_\alpha F \cap \Omega_K$, proving Pieri's formula.

More explicitly, one may construct a sequence of parameterized rational curves $\phi_i : \mathbf{P}^1 \rightarrow \overline{\mathcal{G}}$ for $1 \leq i \leq b$ witnessing this rational equivalence. For $2 \leq i \leq b$, select subspaces $M_i \in U_i$ and pencils $L_{i,t}$ of hyperplanes of M_i by downward induction on i as follows: Choose $M_b \in U_b$. Given $M_i \in U_i$, let $L_{i,t}$ be a pencil of hyperplanes of M_i as in Theorem 3.5, let $M_{i-1} := L_{i,\infty}$, and continue. Then for each i , if $t \neq 0$, $L_{i,t} \in U_{i-1}$. Define $\Sigma_i \subset \mathbf{P}^1 \times \mathbf{G}_m V$ to be the family whose fibre over $t \in \mathbf{P}^1 - \{0\}$ is the variety $Y_{\alpha,i-1}(F, L_{i,t})$.

Let $\psi : \mathbf{P}^1 \rightarrow \overline{U_0} = \mathbf{G}_{n+1-m-s} V$ be a map with $\psi(0) = M_1 := L_{2,\infty}$, $\psi(\infty) = K$, and $\psi^{-1}(U_0) = \mathbf{P}^1 - \{0\}$. Let $\Sigma_1 \subset \mathbf{P}^1 \times \mathbf{G}_m V$ be the family whose fibre over $t \in \mathbf{P}^1$ is $\Omega_\alpha F \cap \Omega_{\psi(t)}$, a generically transverse intersection which is irreducible for $t \neq 0$, by Theorem 2.3. Then for $1 \leq i \leq b$, $\Sigma_i \subset \mathbf{P}^1 \times \mathbf{G}_m V$ is a family with equidimensional generically reduced fibres over $\mathbf{P}^1$.

For $1 \leq i \leq b$, let $\phi_i : \mathbf{P}^1 \rightarrow \overline{\mathcal{G}_{i-1}}$ be the map associated to the family Σ_i , as in §3.1. Then $\phi_i(0) = \phi_{i+1}(\infty) \in \mathcal{G}_i$ and $\phi_i(t) \in \mathcal{G}_{i-1}$ for $t \neq 0$, by Theorem 3.5. Thus these parameterized rational curves give a chain of rational equivalences between $\Omega_\alpha F \cap \Omega_K$ and $Y_{\alpha,b} F$. $\blacksquare$

Let $\beta \in \alpha * r$ and $\gamma \in \alpha * (r+1)$. If $\gamma \in \beta * 1$ with $j(\alpha, \gamma) = j(\beta, \gamma)$, write $\beta \prec_\alpha \gamma$.

3.8. Proof of Theorem 3.5. Let $t \neq 0$. Recall that L_t contains the subspace F_{α_1+s} of M , but not F_{α_1+s-1} . Since $M \in U_{\alpha,s-1} F$, we have $F_{\alpha_1} \cap M = F_{\alpha_1+s-1}$, but $F_{\alpha_1} \cap L_t = F_{\alpha_1+s}$, thus $F_i \cap L_t$ is a hyperplane of $F_i \cap M$ for any $i \leq \alpha_1$. Then $L_t \in U_{\alpha,s} F$, for $t \neq 0$, as

1. $F_{\alpha_1} \cap L_t = F_{\alpha_1+s}$.
2. For $1 \leq j \leq m$, $F_{\alpha_j} \cap M = F_{\alpha_j+1} \cap M$. So $F_{\alpha_j} \cap L_t = F_{\alpha_j+1} \cap L_t$. Moreover, $\dim F_{\alpha_j} \cap L_t = \dim F_{\alpha_j} \cap M - 1$, which is $n + 2 - \alpha_j - j - s$.

Suppose $t \neq 0$ and recall that

$$Y_{\alpha,r}(F, L_t) = \sum_{\substack{\beta \in \alpha * r \\ j(\alpha, \beta) = 1}} \Omega_{\beta+(s-1)\delta^1} F + \sum_{\substack{\beta \in \alpha * r \\ j(\alpha, \beta) > 1}} X_\beta(j(\alpha, \beta), F, L_t).$$

This defines a family $\Sigma \subset (\mathbf{P}^1 - \{0\}) \times \mathbf{G}_m V$ with equidimensional (actually isomorphic) fibres over $\mathbf{P}^1 - \{0\}$. We establish Theorem 3.5, showing the fibre of Σ at 0 is $Y_{\alpha,r+1}(F, M)$ by examining each component of $Y_{\alpha,r}(F, L_t)$ separately, then assembling the result.

Let $\beta \in \alpha * r$. Consider a component of $Y_{\alpha,r}(F, L_t)$ in the first summand, so $j(\alpha, \beta) = 1$. Then $\gamma := \beta + \delta^1$ is the unique sequence satisfying $\beta \prec_\alpha \gamma$. In this case, $\Omega_{\beta+(s-1)\delta^1} F = \Omega_{\gamma+(s-2)\delta^1} F$.

Now consider a component in the second sum, so $j = j(\alpha, \beta) > 1$. Let $\beta' := \beta|_j$, $F' := F|_{\beta_j}$, and $L'_t := F_{\beta_j} \cap L_t$. For $t \neq 0$, Corollary 2.6 gives

$$X_\beta(j(\alpha, \beta), F, L_t) = p(\pi^{-1}(\Omega_{\beta'} F' \cap \Omega_{L'_t})).$$

As $\alpha_j < \beta_j < \alpha_{j-1}$, $\dim L'_t = \dim F_{\beta_j} + 1 - j - (s-1)$, by formula (3.2). For $1 \leq i < j$, $\beta_i = \alpha_i$ and so $\dim L'_t \cap F_{\beta_i} = n + 2 - \beta_i - i - s$. Thus, by Theorem 2.3(3), $\Omega_{\beta'} F'_\bullet \cap \Omega_{L'_t}$ is generically transverse and irreducible. We study the ‘limit’ of these cycles as $t \rightarrow 0$, in the sense of §3.1. Define $L' := \lim_{t \rightarrow 0} L'_t = \lim_{t \rightarrow 0} F_{\beta_j} \cap L_t$, which is $F_{\beta_{j+1}} \cap M$, by Lemma 3.4. Then

- (1) $F_{\alpha_1} \cap L' = F_{\alpha_1} \cap M = F_{\alpha_1+s-1}$.
- (2) For $1 \leq i \leq j$, $F_{\beta_i} \cap L' = F_{\beta_{i+1}} \cap L'$. This follows for $i = j$ because we have $L' \subset F_{\beta_{j+1}} \subset F_{\beta_j}$ and for $i < j$, because $\beta_i = \alpha_i$ and $F_{\alpha_i} \cap M = F_{\alpha_{i+1}} \cap M$. Moreover, for $1 \leq i \leq j$, $\dim F_{\beta_i} \cap L' = n + 2 - \beta_i - i - (s-1)$.

Thus $L' \in U_{\beta', s-1} F'_\bullet$ so $\Omega_{\beta'} F'_\bullet \cap \Omega_{L'}$ is generically transverse, by Theorem 2.3(1). So,

$$\lim_{t \rightarrow 0} X_\beta(j(\alpha, \beta), F_\bullet, L_t) = p(\pi^{-1}(\Omega_{\beta'} F'_\bullet \cap \Omega_{L'})).$$

But $\langle F_{\beta_{i-1}}, F_{\beta_i} \cap L \rangle \subset F_{\beta_{i+1}}$, since $L' \in U_{\beta', s-1} F'_\bullet$. By Remark 2.4,

$$\Omega_{\beta'} F'_\bullet \cap \Omega_{L'} = \sum_{\substack{\gamma' \in \beta' * 1 \\ j(\beta', \gamma') = 1}} \Omega_{\gamma' + (s-2)\delta^1} F'_\bullet + \sum_{\substack{\gamma' \in \beta' * 1 \\ j(\beta', \gamma') > 1}} X_{\gamma'}(j(\beta', \gamma'), F'_\bullet, L').$$

And so $\lim_{t \rightarrow 0} X_\beta(j(\alpha, \beta), F_\bullet, L_t)$ is the cycle

$$\sum_{\substack{\gamma' \in \beta' * 1 \\ j(\beta', \gamma') = 1}} p(\pi^{-1}(\Omega_{\gamma' + (s-2)\delta^1} F'_\bullet)) + \sum_{\substack{\gamma' \in \beta' * 1 \\ j(\beta', \gamma') > 1}} p(\pi^{-1}(X_{\gamma'}(j(\beta', \gamma'), F'_\bullet, L'))).$$

We simplify this expression, beginning with the first sum. Let $\gamma' \in \beta' * 1$ satisfy $j(\beta', \gamma') = 1$. Then by Lemma 2.5, $p(\pi^{-1}(\Omega_{\gamma' + (s-2)\delta^1} F'_\bullet))$ equals $\Omega_{\gamma + (s-2)\delta^1} F_\bullet$, where $\gamma := \beta + \delta^1$ is the unique sequence with $\beta \prec_\alpha \gamma$ and $j(\alpha, \gamma) = 1$.

Consider terms in the second sum, those for which $\gamma' \in \beta' * 1$ with $j(\beta', \gamma') > 1$. Then $p(\pi^{-1}(X_{\gamma'}(j(\beta', \gamma'), F'_\bullet, L')))$ is the subvariety of $\Omega_{\beta'} F'_\bullet$ consisting of those H such that there exists $K \subset H$ with $\dim K = j$, $K \in \Omega_{\gamma'} F'_\bullet$, and $\dim K \cap F'_{\gamma_{j(\beta', \gamma')}} \cap L' \geq 1$.

Let $\gamma := \beta + \delta^{j(\beta', \gamma')}$, the unique sequence with $\beta \prec_\alpha \gamma$ and $j(\alpha, \gamma) = j(\beta', \gamma')$. Then, as $\gamma_{j(\alpha, \gamma)} > \beta_j$, the definition of $F'_\bullet$ implies $F'_{\gamma_{j(\beta', \gamma')}} = F_{j(\alpha, \gamma)} \subset F_{\beta_{j+1}}$. Since $L' = F_{\beta_{j+1}} \cap M$, we see that $F'_{\gamma_{j(\beta', \gamma')}} \cap L' = F_{\gamma_{j(\alpha, \gamma)}} \cap M$. Thus if

$$H \in p(\pi^{-1}(X_{\gamma'}(j(\beta', \gamma'), F'_\bullet, L'))),$$

then $H \in \Omega_\gamma F_\bullet$ and $\dim H \cap F_{\gamma_{j(\alpha, \gamma)}} \cap M \geq 1$, so $H \in X_\gamma(j(\alpha, \gamma), F_\bullet, M)$. The reverse inclusion,

$$X_\gamma(j(\alpha, \gamma), F_\bullet, M) \subset p(\pi^{-1}(X_{\gamma'}(j(\beta', \gamma'), F'_\bullet, L'))),$$

is similar.

This shows that $\lim_{t \rightarrow 0} X_\beta(j(\alpha, \beta), F_\bullet, L_t)$ is the cycle

$$(3.8) \quad \sum_{\substack{\beta \prec_\alpha \gamma \\ j(\alpha, \gamma) = 1}} \Omega_{\gamma + (s-2)\delta^1} F_\bullet + \sum_{\substack{\beta \prec_\alpha \gamma \\ j(\alpha, \gamma) > 1}} X_\gamma(j(\alpha, \gamma), F_\bullet, L).$$

The sets $\{\gamma \mid \beta \prec_\alpha \gamma\}$ for $\beta \in \alpha * r$ partition the set $\alpha * (r+1)$. Thus

$$\lim_{t \rightarrow 0} Y_{\alpha, r}(F_\bullet, L_t) = \sum_{\substack{\gamma \in \alpha * (r+1) \\ j(\alpha, \gamma) = 1}} \Omega_{\gamma + (s-2)\delta^1} F_\bullet + \sum_{\substack{\gamma \in \alpha * (r+1) \\ j(\alpha, \gamma) > 1}} X_\beta(j(\alpha, \gamma), F_\bullet, M),$$

which is $Y_{\alpha, r+1}(F_\bullet, M)$. $\blacksquare$

4. LINK TO SCHENSTED INSERTION

The set $\binom{[n]}{m}$ has a partial order, called the Bruhat order: $\alpha \leq \beta$ if and only if $\Omega_\beta F_\bullet \subset \Omega_\alpha F_\bullet$. Combinatorially, this is $\alpha \leq \beta$ if $\alpha_i \leq \beta_i$ for $1 \leq i \leq m$.

We interpret the behavior of the components $X_\beta(j(\alpha, \beta) F_\bullet, L)$ of the intermediate varieties $Y_{\alpha, i-1}(F_\bullet, L)$ in our proof of Pieri's formula (§3.7) as the branching of a certain subtree of $\binom{[n]}{m}$ with root α . This tree arises similarly in the combinatorial proof of Pieri's formula for Schur polynomials using Schensted insertion given in [F]. To simplify this discussion, assume further that $n > \alpha_1 + b$.

Each rational equivalence of §3.7 is induced by a family Σ_i over $\mathbf{P}^1$ with generic fibre in $\mathcal{G}_{i-1}$ and special fibre in $\mathcal{G}_i$. The components of cycles in $\mathcal{G}_{i-1}$ are indexed by $\beta \in \alpha * (i-1)$, with β th component $\Omega_{\beta + (b+1-i)\delta^1} F_\bullet$, if $j(\alpha, \beta) = 1$, and $X_\beta(j(\alpha, \beta), F_\bullet, L)$ otherwise. In passing to $\mathcal{G}_i$ via ϕ_i , the component $\Omega_{\beta + (b+1-i)\delta^1} F_\bullet$ is unchanged, but reindexed: $\Omega_{\gamma + (b-i)\delta^1} F_\bullet$, where $\gamma := \beta + \delta^1$ is the unique sequence with $\beta \prec_\alpha \gamma$. By equation (3.8), the other components become

$$\sum_{\substack{\beta \prec_\alpha \gamma \\ j(\alpha, \gamma) = 1}} \Omega_{\gamma + (b-i)\delta^1} F_\bullet + \sum_{\substack{\beta \prec_\alpha \gamma \\ j(\alpha, \gamma) > 1}} X_\gamma(j(\alpha, \gamma), F_\bullet, M_i).$$

Thus the component of the generic fibre of Σ_i indexed by $\beta \in \alpha * (i-1)$ becomes a sum of components indexed by $\{\gamma \mid \beta \prec_\alpha \gamma\}$ at the special fibre.

This suggests defining a tree $\mathcal{T}_{\alpha, b}$ whose branching represents the 'branching' of components of $Y_{\alpha, i-1}(F_\bullet, L)$ in these deformations. Let $\mathcal{T}_{\alpha, b} \subset \binom{[n]}{m}$ be the tree with vertex set $\bigcup \{\alpha * i \mid 0 \leq i \leq b\}$ and covering relation $\beta \prec_\alpha \gamma$. This is a tree as $\alpha * i$ is partitioned by the sets $\{\gamma \mid \beta \prec_\alpha \gamma\}$ for $\beta \in \alpha * (i-1)$.

For a decreasing m -sequence α , let $\lambda(\alpha)$ be the partition $(\alpha_1 - m, \alpha_2 - m + 1, \dots, \alpha_m - 1)$. The association $\alpha \longleftrightarrow \lambda(\alpha)$ gives an order isomorphism between the set of decreasing m -sequences and the set of partitions of length at most m . This transfers notions for sequences into corresponding notions for partitions.

To a (semi-standard) Young tableau T with entries among $1, \dots, m$, associate a monomial x^T in the variables $x_1, x_2, \dots, x_m$: The exponent of x_i in x^T is the number of occurrences of i in T . The Schur polynomial s_λ is $\sum x^T$, the sum over all tableaux T of shape λ . There is surjective homomorphism from the algebra of Schur polynomials to the Chow ring of $\mathbf{G}_m V$ defined by:

$$s_\lambda \longmapsto \begin{cases} [\Omega_\alpha F_\bullet] & \text{if } \lambda = \lambda(\alpha) \text{ for some } \alpha \in \binom{[n]}{m} \\ 0 & \text{otherwise} \end{cases}.$$

Special Schur polynomials are indexed by partitions $(b, 0, \dots, 0)$ with a single row.

Schensted insertion gives a combinatorial proof of Pieri's formula, providing a content-preserving bijection between the set of pairs (S, T) of tableaux where S has shape λ and T has shape $(b, 0, \dots, 0)$ and the set of all tableaux whose shape is in $\lambda * b$: Insert the reading word of T into S . The resulting tableau has shape $\mu \in \lambda * b$.

Let $\lambda = \lambda_0, \lambda_1, \dots, \lambda_b = \mu$ be the sequence of shapes resulting from the insertion of successive entries of T into S . Since T is a single row, it is a property of the insertion algorithm that $\lambda_i \prec_\lambda \lambda_{i+1}$, and so this sequence is a chain in the tree $\mathcal{T}_{\alpha, b}$.

The totality of these insertions for all such pairs of tableaux gives all chains in $\mathcal{T}_{\alpha, b}$. Thus the 'branching' of shapes during Schensted insertion is identical to the branching of components in the rational equivalences of §3.7. We feel this relation to combinatorics is one of the more intriguing aspects of our proof of Pieri's formula. It leads us to speculate that similar ideas may yield a geometric proof of the Littlewood-Richardson rule.

REFERENCES

- [BS] N. BERGERON AND F. SOTTILE, *Symmetries of Littlewood-Richardson coefficients for Schubert polynomials*, in preparation.
- [HB] B. BOE AND H. HILLER, *Pieri formula for SO_{2n+1}/U_n and Sp_n/U_n* , Adv. in Math., 62 (1986), pp. 49–67.
- [C] C. CHEVALLEY, *Sur les décompositions cellulaires des espaces G/B* , in Algebraic Groups and their Generalizations: Classical Methods, American Mathematical Society, 1994, pp. 1–23. Proceedings and Symposia in Pure Mathematics, vol. 56, Part 1.
- [F] W. FULTON, *Young Tableaux, with Applications to Representation Theory and Geometry*, Cambridge University Press, 1996.
- [H] A. HIRSCHOWITZ, *Le groupe de Chow équivariant*, C. R. Acad. Sci. Paris, 298 (1984), pp. 87–89.
- [HP] W. V. D. HODGE AND D. PEDOE, *Methods of Algebraic Geometry*, vol. II, Cambridge University Press, 1952.
- [K] S. KLEIMAN, *The transversality of a general translate*, Comp. Math., 28 (1974), pp. 287–297.
- [P] P. PRAGACZ, *Algebro-geometric applications of Schur s - and q -polynomials*, in Topics in Invariant Theory, Séminaire d'Algèbre Dubreil-Malliavin 1989-90, Springer-Verlag, 1991, pp. 130–191.
- [PR] P. PRAGACZ AND J. RATAJSKI, *Pieri-type formula for $Sp(2m)/p$ and $SO(2m+1)/p$* , C. R. Acad. Sci. Paris, 317 (1993), pp. 1035–1040.
- [Sa] P. SAMUEL, *Méthodes d'Algèbre Abstraite en Géométrie Algébrique*, Ergebnisse der Math., Springer-Verlag, seconde ed., 1967.
- [Sc] C. SCHENSTED, *Longest increasing and decreasing subsequence*, Canad. J. Math., 13 (1961), pp. 179–191.
- [So1] F. SOTTILE, *Enumerative geometry for the real Grassmannian of lines in projective space*, Duke Math. J., to appear.
- [So2] ———, *Pieri's formula for flag manifolds and Schubert polynomials*, Annales d'Institut Fourier, Grenoble, 46 (1996), pp. 1–22.
- [St] J. STEMBRIDGE, *Shifted tableaux and the projective representations of the symmetric group*, Adv. Math., 74 (1989), pp. 87–134.

APPENDICES

Following are two appendices which are for general distribution and not intended for publication. In the first, we illustrate how one may introduce coordinates by giving an explicit parameterized family of intersections in one example and studying how the intersection cycles deform. It should be possible to transfer this description into a parameterized family of ideals, and use computer algebra to study the deformations.

In the second, we illustrate how deformations of the kind studied here may be used to find real solutions to some problems of enumerative geometry.

APPENDIX A: AN EXAMPLE ILLUSTRATING EXPLICIT RATIONAL EQUIVALENCE

Our proof of Pieri's formula used methods which were almost¹ entirely synthetic (coordinate-free). Here, we illustrate our methods on a particular example, simultaneously showing how to introduce coordinates. While we do not attempt such an analysis, it should be possible to replicate our results using (computer) algebra. It is in this context that these 'explicit rational equivalences' are most explicit.

A.1 Example: $n = 9$, $m = 3$, $s = 2$, and $\alpha = 741$. Let F be a flag and L a five-dimensional subspace of $V = k^9$ which meets F properly. We choose coordinates for V and give a two parameter family of 5-planes containing L which exhibits a 'chain of rational equivalences' between $\Omega_{741}F \cap \Omega_L$ and $Y_{741,2}F = \sum_{\beta \in 741*2} \Omega_{\beta}F$, which is the cycle

$$\Omega_{941}F + \Omega_{851}F + \Omega_{761}F + \Omega_{842}F + \Omega_{752}F + \Omega_{743}F.$$

Let $e_1, \dots, e_9$ be a basis for V such that $F_j = \langle e_j, \dots, e_9 \rangle$, for $1 \leq j \leq 9$. Let $x_1, \dots, x_9$ be dual coordinates for V^* , then F_j is defined by the linear forms $x_1, \dots, x_{j-1}$. Further assume that $e_1, \dots, e_9$ have been chosen so that $L = \langle e_1, e_2, e_3, e_4, e_5 \rangle$.

For $(s, t) \in k^2$, define linearly independent forms $\Lambda_1, \Lambda_2, \Lambda_3$, and Λ_4 in V^* :

$$\begin{aligned} \Lambda_1 &:= x_1 + sx_8 \\ \Lambda_2 &:= x_2 + tx_3 + st^2x_4 + (t^2 + st^3)x_5 + (t^3 + st^4)x_6 + t^4x_8 \\ \Lambda_3 &:= x_4 + sx_9 \\ \Lambda_4 &:= x_7 \end{aligned}$$

If $s \cdot t \neq 0$, let $\sigma = 1/s$ and $\tau = 1/t$. Consider the forms

$$\begin{aligned} \sigma\tau^4\Lambda_2 - \sigma^2\Lambda_1 &= -\sigma^2x_1 + \sigma\tau^4x_2 + \sigma\tau^3x_3 + \tau^2x_4 + (\sigma\tau^2 + \tau)x_5 + (\sigma\tau + 1)x_6 \\ \Lambda_4 &= x_7 \\ \sigma\Lambda_1 &= \sigma x_1 + x_8 \\ \sigma\Lambda_3 &= \sigma x_4 + x_9. \end{aligned}$$

These are linearly independent for $(\sigma, \tau) \in k^2$. These two sets of forms together define a family $L_{s,t}$ of 5-dimensional subspaces of k^9 with base $U := \mathbf{P}^1 \times \mathbf{P}^1 - \{(\infty, 0), (0, \infty)\}$, and with $L_{\infty, \infty} = L$. For $s \cdot t \neq 0$, the second set of forms shows that $L_{s,t}$ meets the subspaces F_1, F_4 , and F_7 properly. This shows that for $s \cdot t \neq 0$ all 5-planes $L_{s,t}$ are in the dense orbit of the parabolic subgroup P stabilizing F_1, F_4 , and F_7 .

¹ Coordinates were only chosen in the proof of Lemma 3.4. A synthetic proof, while possible, is perhaps less convincing, and certainly not as brief.

Thus over $(\mathbf{P}^1 - \{0\}) \times (\mathbf{P}^1 - \{0\})$, there is a family Σ of cycles with fibre over (s, t) :

$$\Sigma_{s,t} := \Omega_{741}F \cdot \bigcap \Omega_{L_{s,t}}. \quad (A.1)$$

Moreover, all fibres of Σ are isomorphic by some element of P . We study the cycles obtained as first $s \rightarrow 0$ with $t \neq 0$ fixed, and then $t \rightarrow 0$. We show:

- (A) For $s \cdot t \neq 0$, the intersection $\Omega_{741}F \cdot \bigcap \Omega_{L_{s,t}}$ is generically transverse and irreducible.
- (B) For $t \neq 0$ or ∞ , the intersection cycle $Y_t := \Omega_{741}F \cdot \bigcap \Omega_{L_{0,t}}$ is generically transverse, but reducible, with components indexed by $\{841, 751, 742\} = 741 * 1$. This is because if $t \neq 0$ or ∞ , then $L_{0,t} \in U_{741,2}F$. Thus Σ can be extended to $U - \{k \times 0\}$ by (§A.1).
- (C) The intersection $\Omega_{741}F \cdot \bigcap \Omega_{L_{0,0}}$ is improper. However, $\lim_{t \rightarrow 0} Y_t = \sum_{\beta \in 741*2} \Omega_{\beta}F$.

We show these points:

(A): We noted already that for $s \cdot t \neq 0$, $L_{s,t}$ meets F_1, F_4 , and F_7 properly. Then by Theorem 2.3, the intersection $\Omega_{741}F \cdot \bigcap \Omega_{L_{s,t}}$ is generically transverse and irreducible.

(B): Fix $t \neq 0, \infty$, and consider the family $L_{s,t}$ for $s \in \mathbf{P}^1$. Then $L_{0,t}$ is given by the forms:

$$x_1, \quad x_4, \quad x_7, \quad x_2 + tx_3 + t^2x_5 + t^3x_6 + t^4x_8$$

and so it does not meet F properly. Note that $L_{0,t}$ has a basis:

$$te_2 - e_3, \quad te_3 - e_5, \quad te_5 - e_6, \quad te_6 - e_8, \quad e_9.$$

Then $L_{0,t} \in U_{741,2}F$ as

1. $L_{0,t} \subset F_2$.
2. $L_{0,t} \cap F_4 = L_{0,t} \cap F_5 = \langle te_5 - e_6, \quad te_6 - e_8, \quad e_9 \rangle$, which has dimension 3.
3. $L_{0,t} \cap F_7 = F_9$.

Note also that $F_2 = \langle L_{0,t}, F_4 \rangle$, $F_5 = \langle L_{0,t} \cap F_4, F_7 \rangle$, and $F_8 \supset \langle L_{0,t} \cap F_7 \rangle$. By Theorem 2.3,

$$\Omega_{741}F \cdot \bigcap \Omega_{L_{0,t}} = X_{841}(1, F, L_{0,t}) + X_{751}(2, F, L_{0,t}) + X_{742}(3, F, L_{0,t}).$$

(C): Note $L_{0,0} = \langle e_3, e_5, e_6, e_8, e_9 \rangle$. Since $F_8 \subset L_{0,0}$, $\Omega_{841}F \cdot \bigcap \Omega_{L_{0,0}}$ is improper.

Instead let Σ be the family of cycles over $\mathbf{P}^1$ whose fibre over $t \neq 0, \infty$ is

$$\Sigma_t := \Omega_{741}F \cdot \bigcap \Omega_{L_{0,t}},$$

and we describe the fibre $\Sigma_0 \subset \Omega_{741}F \cdot \bigcap \Omega_{L_{0,0}}$. For this, we need a better description of the fibres Σ_t , for $t \neq 0, \infty$. First, let $M = \langle e_2, e_3, e_5, e_6, e_8, e_9 \rangle \in U_{741,1}F$.

- $j = 1$: $X_{841}(1, F, L_{0,t}) = \{H \in \Omega_{841}F \mid \dim H \cap F_8 \cap L_{0,t} \geq 1\}$. Since $L_{0,t} \cap F_8 = F_9$, this equals $\Omega_{941}F = \Omega_{841+100}F$.
- $j = 2$: Similarly, $X_{751}(2, F, L_{0,t}) = \{H \in \Omega_{751}F \mid \dim H \cap F_5 \cap L_{0,t} \geq 1\}$. If p, π are the first and second projections on

$$\widetilde{\Omega_{751}F}^2 = \{(H, K) \in \Omega_{751}F \times \mathbf{G}_3F_5 \mid K \subset H, \dim K \cap F_7 \geq 1\},$$

then $X_{751}(2, F, L_{0,t}) = p(\pi^{-1}(\Omega_{31}F|_5 \bigcap \Omega_{F_5 \cap L_{0,t}}))$.

Since $F_5 \cap L_{0,t} = \langle e_6 - te_5, e_8 - te_6, e_9 \rangle$, $\lim_{t \rightarrow 0} F_5 \cap L_{0,t} = F_6 \cap M$.

Suppose $K \in \Omega_{31}F_\bullet|_5 \cap \Omega_{F_6 \cap M}$. Then $K \subset F_5$, $\dim K \cap F_7 \geq 1$, and $\dim K \cap F_6 \cap M \geq 1$.

- (a) If $K \cap (F_6 \cap M) \subset F_7$, then $\dim K \cap F_8 \geq 1$, as $F_6 \cap M \cap F_7 = F_8$, and so $K \in \Omega_{41}F_\bullet|_5$. Also, $p(\pi^{-1}(\Omega_{41}F_\bullet|_5)) = \Omega_{851}F_\bullet$.
- (b) If $K \cap F_6 \cap M \not\subset F_7$, then $K \subset \langle F_7, M \cap F_6 \rangle = F_6$ and so $K \in \Omega_{32}F_\bullet|_5$. Note that $p(\pi^{-1}(\Omega_{32}F_\bullet|_5)) = \Omega_{761}F_\bullet$.

This shows that $\Omega_{31}F_\bullet|_5 \cap \Omega_{F_6 \cap M} = \Omega_{41}F_\bullet|_5 + \Omega_{32}F_\bullet|_5$. Since this intersection is generically transverse, it follows that

$$\lim_{t \rightarrow 0} X_{751}(2, F_\bullet, L_{0,t}) = p(\pi^{-1}(\Omega_{41}F_\bullet|_5 + \Omega_{21}F_\bullet|_5)) = \Omega_{851}F_\bullet + \Omega_{761}F_\bullet.$$

$j = 3$: $X_{742}(3, F_\bullet, L_{0,t}) = \{H \in \Omega_{742}F_\bullet \mid \dim H \cap L_{0,t} \cap F_2 \geq 1\}$. This is $\Omega_{631}F_\bullet|_2 \cap \Omega_{L_{0,t}}$, which is generically transverse and irreducible, *as an intersection in $\mathbf{G}_3F_2$* , by Theorem 2.3.

Similarly, $X_{742}(3, F_\bullet, L_{0,0}) = \Omega_{631}F_\bullet|_2 \cap \Omega_{L_{0,0}}$ is generically transverse but reducible, as an intersection in $\mathbf{G}_3F_2$, by Theorem 2.3(2). Thus

$$\lim_{t \rightarrow 0} X_{742}(3, F_\bullet, L_{0,t}) = X_{742}(3, F_\bullet, L_{0,0}).$$

Let $H \in X_{742}(3, F_\bullet, L_{0,0})$. Then $H \subset F_2$, $\dim H \cap F_4 \geq 2$, $\dim H \cap F_7 \geq 1$, and $\dim H \cap L_{0,0} \geq 1$.

- (a) If $H \cap L_{0,0} \subset F_7$, then $\dim H \cap F_8 \geq 1$, as $F_7 \cap L_{0,0} = F_8$. Thus $H \in \Omega_{842}F_\bullet$.
- (b) If $\dim H \cap F_4 \cap L_{0,0} \geq 1$ and $H \cap L_{0,0} \not\subset F_7$, then $\dim H \cap \langle F_7, F_4 \cap L_{0,0} \rangle \geq 2$. Since $F_5 = \langle F_7, F_4 \cap L_{0,0} \rangle$, $H \in \Omega_{752}F_\bullet$.
- (c) If $H \cap L_{0,0} \not\subset F_4$, then $H \subset \langle F_4, F_2 \cap L_{0,0} \rangle = F_3$, and so $H \in \Omega_{743}F_\bullet$.

We deduce that $X_{742}(3, F_\bullet, L_{0,0}) = \Omega_{842}F_\bullet + \Omega_{752}F_\bullet + \Omega_{743}F_\bullet$, and so

$$\lim_{t \rightarrow 0} X_{742}(3, F_\bullet, L_{0,t}) = \Omega_{842}F_\bullet + \Omega_{752}F_\bullet + \Omega_{743}F_\bullet.$$

These calculations identify the fibre Σ_0 of the family whose fibre over $t \neq 0$ is $\Omega_\alpha F_\bullet \cap \Omega_{L_{0,t}}$. They show $\Sigma_0 = \Omega_{841+100}F_\bullet + X_{751}(2, F_\bullet, L_{0,0}) + X_{742}(3, F_\bullet, L_{0,0})$, which is


$$\Omega_{941}F_\bullet + \Omega_{851}F_\bullet + \Omega_{761}F_\bullet + \Omega_{842}F_\bullet + \Omega_{752}F_\bullet + \Omega_{743}F_\bullet = \sum_{\beta \in 741*2} \Omega_\beta F_\bullet.$$

A.2: The rational equivalences, explicitly The calculations of §A.1 yield an explicit chain of rational equivalences from $\Omega_{741}F_\bullet \cap \Omega_L$ to $\sum_{\beta \in 741*2} \Omega_\beta F_\bullet$. Furthermore, the generic fibre of Σ_1 is irreducible and the special fibre has three components, indexed by the set $741*1$. The generic fibre of Σ_2 also has three components indexed by $741*1$, and its special fibre has six components, indexed by the set $741*2$. The tree $\mathcal{T}_{741,2}$, shown in Figure 1 in the lattice of Young diagrams, represents this branching of components.

APPENDIX B: APPLICATION TO REAL ENUMERATIVE GEOMETRY

As in [So1], explicit rational equivalences can be used to find real solutions to problems in enumerative geometry. Suppose $k = \mathbf{R}$ and $F_\bullet$ is a real flag.

Lemma B.1 *Let $F_\bullet$ be a flag in $\mathbf{R}^n$, $\alpha \in \binom{[n]}{m}$, and $b \geq 0$. Suppose X is a subvariety of $\mathbf{G}_m V$ which meets $Y_{\alpha,b}F_\bullet$ transversally in d points, with e real. Then there exists $L \in \mathbf{G}_{n+1-m-b}\mathbf{R}^n$ with $\Omega_\alpha F_\bullet \cap \Omega_L$ a generically transverse and irreducible intersection such that X meets $\Omega_\alpha F_\bullet \cap \Omega_L$ transversally in d points, with e real.*

FIGURE 1. The Tree $\mathcal{T}_{741,2}$

Proof: The key idea is that the number of real points in the fibre of a finite real morphism is preserved under real deformations.

We use the notation of §3.7. For $1 \leq i \leq b$, let $\mathcal{G}_{i;\mathbf{R}} \subset \mathcal{G}_i$ be those cycles $Y_{\alpha,i}(F, L)$ with L a real subspace. Similarly, let $\mathcal{G}_{0;\mathbf{R}} \subset \mathcal{G}_0$ be the cycles $\Omega_{\alpha}F \cap \Omega_L$ with L real. The description of the maps ϕ_i show they are real algebraic.

Suppose $(\Sigma_i)_0 = Y_{\alpha,i}(F, M_i)$ meets X transversally in d points, with e real. For t in $\mathbf{RP}^1 - \{0\}$, $(\Sigma_i)_t \in \mathcal{G}_{i-1;\mathbf{R}}$, since Σ_i is induced by a real pencil $L_{i,t}$. Since $(\Sigma_i)_0$ meets X transversally in d points, there is a Zariski open subset U of $\mathbf{P}^1$ containing 0 over which each fibre $(\Sigma_i)_t$ of Σ_i meets X transversally in d points. Let I be the connected component of $U(\mathbf{R})$ containing 0. For $t \in I$, the fibres $(\Sigma_i)_t$ meet X transversally in d points, with e real, as the number of real points in the intersection is constant on connected components of $U(\mathbf{R})$.

Downward induction on i completes the proof, where, to continue, either reparameterize $\mathbf{P}^1$ so $\infty \in I$, or choose M_{i-1} to be $L_{i,t}$ for some $t \in I - \{0\}$. $\blacksquare$

For $\alpha \in \binom{[n]}{m}$, let $\alpha^\vee \in \binom{[n]}{m}$ be $n+1-\alpha_m > \dots > n+1-\alpha_1$. Then α and $\alpha^\vee$ are the indices of Poincaré dual classes.

Theorem B.2 Let $\alpha, \beta \in \binom{[n]}{m}$ and a, b, c be positive integers with $a + b + c + |\alpha| + |\beta| = m(n-m)$, the dimension of $\mathbf{G}_m \mathbf{R}^n$. Let d be the number of pairs $(\gamma, \delta) \in (\alpha * a) \times (\beta * b)$ satisfying $\delta^\vee \in \gamma * c$. Then there exist flags $F_\bullet$ and $F'_\bullet$ and subspaces A, B , and C of $\mathbf{R}^n$ of respective dimensions $n+1-m-a, n+1-m-b$, and $n+1-m-c$ such that $\Omega_\alpha F_\bullet, \Omega_\beta F'_\bullet, \Omega_A, \Omega_B$, and Ω_C meet transversally in d real points.

The conditions on α, β, a, b, c ensure that for general F, F', A, B , and C , the Schubert varieties meet in a degree d zero cycle. In fact, we prove this below.

Proof: First suppose $a = b = 0$ so $\Omega_A = \Omega_B = \mathbf{G}_m \mathbf{R}^n$. This triple intersection arises in the classical proof of Pieri's formula (*cf.* [HP]): Suppose F, F' , and C are in linear general position and $c + |\alpha| + |\beta| = m(n - m)$. If $\beta^\vee \in \alpha * c$, the varieties $\Omega_\alpha F$, $\Omega_\beta F'$, and Ω_C meet transversally in a single m -plane $H_{\alpha\beta}$, otherwise their intersection is empty.

Let $\beta^\vee \in \alpha * c$. Then $H_{\alpha\beta}$ has a basis $f_1, \dots, f_m$ with $f_j \in K_j := F_{\alpha_j} \cap F'_{\beta_{m+1-j}}$, where $f_1 + \dots + f_m$ spans the line $C \cap (K_1 \oplus \dots \oplus K_m)$. The conditions on α, β , and c ensure that when F, F' , and C are in linear general position, each $K_i \neq \{0\}$, the sum $K_1 + \dots + K_m$ is direct, the dimension of $C \cap (K_1 \oplus \dots \oplus K_m)$ is 1, that $f_i \notin F_{\alpha_i+1}$, and that $f_i \notin F'_{\beta_{m+1-i}+1}$. Thus $H_{\alpha\beta}$ is real and if $(\alpha, \beta) \neq (\alpha', \beta')$, then $H_{\alpha\beta} \neq H_{\alpha'\beta'}$.

We return to the general case. The previous discussion shows that if F, F' , and C are real subspaces in linear general position, then

$$\left(\sum_{\gamma \in \alpha * a} \Omega_\gamma F \right) \cap \left(\sum_{\delta \in \beta * b} \Omega_\delta F' \right) \cap \Omega_C$$

is a transverse intersection consisting of the d real m -planes:

$$\{H_{\gamma\delta} \mid \gamma \in \alpha * a, \delta \in \beta * b, \text{ and } \delta^\vee \in \gamma * c\}.$$

By Lemma B.1, with $Y = \left(\sum_{\delta \in \beta * b} \Omega_\delta F' \right) \cap \Omega_C$, there exists $A \in \mathbf{G}_{n+1-m-a} \mathbf{R}^n$ with

$$\Omega_\alpha F \cap \Omega_A \cap \left(\sum_{\delta \in \beta * b} \Omega_\delta F' \right) \cap \Omega_C$$

a transverse intersection consisting of d real m -planes. Another application of Lemma B.1 with $Y = \Omega_\alpha F \cap \Omega_A \cap \Omega_C$ shows there exists $B \in \mathbf{G}_{n+1-m-a} \mathbf{R}^n$ with

$$\Omega_\alpha F \cap \Omega_\beta F' \cap \Omega_A \cap \Omega_B \cap \Omega_C$$

a transverse intersection consisting of d real m -planes. $\blacksquare$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 100 ST. GEORGE STREET, TORONTO, ONTARIO M5S 3G3, CANADA, (416) 978-4031

E-mail address: sottle@math.toronto.edu